

THE PSCZ GALAXY POWER SPECTRUM COMPARED TO N-BODY SIMULATIONS

Mark C. Neyrinck, Andrew J.S. Hamilton, Nickolay Y. Gnedin
JILA and Dept. of Astrophysical & Planetary Sciences, University of Colorado
 Mark.Neyrinck@colorado.edu, Andrew.Hamilton@colorado.edu, gnedin@casa.colorado.edu

Abstract By comparing the PSCz galaxy power spectrum with haloes from nested and phased N-body simulations, we try to understand how IRAS infrared-selected galaxies populate dark-matter haloes. We pay special attention to the way we identify haloes in the simulations.

The galaxy power spectrum measured by Hamilton & Tegmark (2002) from the PSCz redshift survey (Saunders et al, 2000) has a remarkable power-law form over 4 decades of wavenumber. Interpreting it in the Λ CDM paradigm requires scale-dependent bias between galaxies and dark matter, which we hoped to reproduce in N-body simulations.

We ran four 256^3 -particle AP^3M dark matter simulations of comoving boxsize 256, 128, 64, and $32 h^{-1}$ Mpc, each with a fixed Eulerian softening length of $10 h^{-1}$ kpc. The boxes are nested in the sense that the phases of initial fluctuations match in the centers of each box, ensuring similar structures there. (For a movie depicting this, look at <http://casa.colorado.edu/~neyrinck/nesthalf.mpg>)

The non-trivial step in this project is to go from a collection of dark matter particles to a set of haloes that could possibly host real galaxies. We used DENMAX (Gelb & Bertschinger 1994) with a smoothing length of one fifth the mean interparticle separation, which gives a mass spectrum similar to that given by Press-Schechter (1974). We then varied a low mass cutoff in the list of haloes to obtain the best fit to PSCz. However, there seemed to be an exclusion effect of close pairs because of the rather large smoothing length, showing itself as a small-scale downturn in the halo power spectra.

We thus tried “DENMAX²,” running DENMAX with half the previous smoothing length on each halo separately to resolve subhaloes. We wanted to characterize sets of DENMAX and DENMAX² haloes with the

same parameter (mass would no longer do, since the mass returned by DENMAX changes with smoothing length), so we used central density, which turns out to be well-correlated with mass in our simulations.

As expected, DENMAX² resolved more close pairs than DENMAX, extending the correlation function power law at the small-scale end by a factor of two or three. Fig. 1 shows the best fits from each of the four nested simulations. The central density cutoff in each simulation corresponded to a mass cutoff of about $10^{11} M_{\odot}$.

In conclusion, we have succeeded in reproducing the scale-dependent bias of the PSCz power spectrum at scales $\gtrsim 30 h^{-1} \text{ kpc}$ with N-body simulations. Extending the analysis to the smallest scales requires resolving the substructure of dark matter haloes, which we have done with an algorithm we call DENMAX².

References

- Gelb J., & Bertschinger, E. 1994, ApJ, 436, 467
 Hamilton, A.J.S., & Tegmark, M. 2002, MNRAS, 330, 506
 Press, W.H., & Schechter, P. 1974, ApJ, 187, 425
 Saunders, W., et al. 2000, MNRAS, 317, 55




Figure 1 The PSCz galaxy power spectrum and its errors, along with best fits from the 256, 128, 64, and 32 $h^{-1} \text{ Mpc}$ simulations (error bars suppressed). Each fit is good over a range of ~ 256 in wavenumber.